It is *Not* Possible to Design a Rotationally Invariant Sampling Scheme for DT-MRI

D. K. Jones[1], C. G. Koay[2], and P. J. Basser[2]

[1]CUBRIC and School of Psychology, Cardiff, Wales, United Kingdom, [2]STBB/LIMB/NICHD, National Institutes of Health, Bethesda, MD, United States

INTRODUCTION: The choice of gradient sampling orientations for DT-MRI experiments has exercised many groups in the field. Of particular interest is the design of sampling schemes that will ensure that the error / variance in tensor estimates is independent of the relative orientation of the tensor to the reference frame established by the sampling vectors, which others have termed 'statistical rotational invariance' or '*SRI*'[1,2]. Previous work has identified a relationship between the condition number of the quadratic encoding matrix, formed from the gradient sampling orientations, and the variance in the estimated diffusion tensor[3]. Batchelor *et al.*[4] showed that certain sampling schemes (e.g., the dual-gradient scheme) had rotationally *variant* condition numbers while, those derived from the vertices of an icosahedron had rotationally *invariant* condition numbers. Thus, schemes based on vectors pointing to the vertices of an icosahedron (or tessellated icosahedrons) cause initial excitement. However, Jones[2] showed that, in discord with this theory, not all icosahedral schemes are the same – and that rotationally invariant condition number is a necessary but insufficient requirement for rotational invariance. A more general framework has been proposed[1,5] – which provides a template for sampling schemes in terms of the fourth order precision matrix. Previous theoretical considerations suggested that, under the *linear* framework, a statistically rotationally invariant sampling scheme could not be designed, except for the trivial case of isotropic tensors[5]. Here, we address an important and long outstanding question in the DT-MRI literature, i.e. In the limit of an infinite number of sampling orientations, can one design a sampling scheme which is statistically rotationally invariant?

THEORY: By combining results from previous works[1,5,6], we can show how the precision matrix, **M'**, in estimation of the diffusion tensor, **D**, can be expressed in terms of the elements of the *m* unit sampling vectors, **g** , the *b*-values, b_m and resultant **B** matrices, $\mathbf{B}_m$ (where $\mathbf{B}_m = b_m \mathbf{g}_m^{\mathrm{T}} \mathbf{g}_m$), via Eq. [1] , where $\alpha = 0$ and 1 for non-linear and linear regression, respectively:

$$\mathbf{M}' = \begin{bmatrix} \sum_m \frac{b_m^2 g_{X_m}^4}{\alpha+\exp(2Tr(\mathbf{B}_m\mathbf{D})} & \sum_m \frac{b_m^2 g_{X_m}^2 g_{Y_m}^2}{\alpha+\exp(2Tr(\mathbf{B}_m\mathbf{D})} & \sum_m \frac{b_m^2 g_{X_m}^2 g_{Z_m}^2}{\alpha+\exp(2Tr(\mathbf{B}_m\mathbf{D})} & 2\sum_m \frac{b_m^2 g_{X_m}^3 g_{Y_m}}{\alpha+\exp(2Tr(\mathbf{B}_m\mathbf{D})} & 2\sum_m \frac{b_m^2 g_{X_m}^3 g_{Z_m}}{\alpha+\exp(2Tr(\mathbf{B}_m\mathbf{D})} & 2\sum_m \frac{b_m^2 g_{X_m} g_{Y_m} g_{Z_m}}{\alpha+\exp(2Tr(\mathbf{B}_m\mathbf{D})} \\ . & \sum_m \frac{b_m^2 g_{Y_m}^4}{\alpha+\exp(2Tr(\mathbf{B}_m\mathbf{D})} & \sum_m \frac{b_m^2 g_{Y_m}^2 g_{Z_m}^2}{\alpha+\exp(2Tr(\mathbf{B}_m\mathbf{D})} & 2\sum_m \frac{b_m^2 g_{X_m} g_{Y_m}^3}{\alpha+\exp(2Tr(\mathbf{B}_m\mathbf{D})} & 2\sum_m \frac{b_m^2 g_{X_m} g_{Y_m}^2 g_{Z_m}}{\alpha+\exp(2Tr(\mathbf{B}_m\mathbf{D})} & 2\sum_m \frac{b_m^2 g_{Y_m}^3 g_{Z_m}}{\alpha+\exp(2Tr(\mathbf{B}_m\mathbf{D})} \\ . & . & \sum_m \frac{b_m^2 g_{Z_m}^4}{\alpha+\exp(2Tr(\mathbf{B}_m\mathbf{D})} & 2\sum_m \frac{b_m^2 g_{X_m} g_{Y_m} g_{Z_m}^2}{\alpha+\exp(2Tr(\mathbf{B}_m\mathbf{D})} & 2\sum_m \frac{b_m^2 g_{X_m} g_{Y_m}^3}{\alpha+\exp(2Tr(\mathbf{B}_m\mathbf{D})} & 2\sum_m \frac{b_m^2 g_{Y_m} g_{Z_m}^3}{\alpha+\exp(2Tr(\mathbf{B}_m\mathbf{D})} \\ . & . & . & 4\sum_m \frac{b_m^2 g_{X_m}^2 g_{Y_m}^2}{\alpha+\exp(2Tr(\mathbf{B}_m\mathbf{D})} & 4\sum_m \frac{b_m^2 g_{X_m}^2 g_{Y_m} g_{Z_m}}{\alpha+\exp(2Tr(\mathbf{B}_m\mathbf{D})} & 4\sum_m \frac{b_m^2 g_{X_m} g_{Y_m}^2 g_{Z_m}}{\alpha+\exp(2Tr(\mathbf{B}_m\mathbf{D})} \\ . & . & . & . & 4\sum_m \frac{b_m^2 g_{X_m}^2 g_{Z_m}^2}{\alpha+\exp(2Tr(\mathbf{B}_m\mathbf{D})} & 4\sum_m \frac{b_m^2 g_{X_m} g_{Y_m} g_{Z_m}^2}{\alpha+\exp(2Tr(\mathbf{B}_m\mathbf{D})} \\ . & . & . & . & . & 4\sum_m \frac{b_m^2 g_{Y_m}^2 g_{Z_m}^2}{\alpha+\exp(2Tr(\mathbf{B}_m\mathbf{D})} \end{bmatrix} \quad [1]$$

It has been shown that for the precision matrix to be statistically rotationally invariant, it should take the following general form[1,7]:

$$\mathbf{M}_{\mathbf{iso}} = \begin{bmatrix} \lambda+2\mu & \lambda & \lambda & 0 & 0 & 0 \\ \lambda & \lambda+2\mu & \lambda & 0 & 0 & 0 \\ \lambda & \lambda & \lambda+2\mu & 0 & 0 & 0 \\ 0 & 0 & 0 & 4\mu & 0 & 0 \\ 0 & 0 & 0 & 0 & 4\mu & 0 \\ 0 & 0 & 0 & 0 & 0 & 4\mu \end{bmatrix} \quad [2]$$

In Eq. [1], $\mathbf{M}'_{4,4} = 4\mathbf{M}'_{1,2}$, $\mathbf{M}'_{5,5} = 4\mathbf{M}'_{1,3}$ and $\mathbf{M}'_{6,6} = 4\mathbf{M}'_{2,3}$, which on comparison with Eq. [2] means that λ must be equal to μ.. To conform to the prescription in Eq. [2], *all* elements of M' should conform for *all* possible tensors. Therefore, if we can find just one example of non-conformance, we have shown that the design is not *SRI*. To make progress, we examine just the non-zero elements of M', which contain only even powers of the elements of sampling vector **g**. Thus, we define an even function of these elements, as: $\kappa(\mathbf{g}_p) = (g_{X_p})^{2l}(g_{Y_p})^{2m}(g_{Z_p})^{2n}$. Summing over an *infinite* number of directions, is the same as integrating over all angles – so we re-parameterize **g** in spherical co-ordinates, **g** = [$\sin\theta\cos\phi$, $\sin\theta\sin\phi$, $\cos\phi$], thus the form for the infinite sums in Eq. [1] becomes

$$< f > = \frac{b^2}{4\pi} \int_{\phi=0}^{2\pi} \int_{\theta=0}^{\pi} \exp(-2Tr(\mathbf{BD}))\kappa(\mathbf{g}) \sin(\theta) d\theta d\phi \quad [3]$$

If we assume **D** is aligned with principal lab frame, (i.e., $\mathbf{D}_{ij}=\delta_{ij}\lambda_i$, where δ_{ij} is the Dirac delta function, and λ_i are eigenvalues), then

$$< f > = \frac{b^2}{4\pi} \int_{\phi=0}^{2\pi} \int_{\theta=0}^{\pi} (g_X)^{2l}(g_Y)^{2m}(g_Z)^{2n} \exp(-(s_1 g_X^2 + s_2 g_Y^2 + s_3 g_Z^2)) \sin(\theta) d\theta d\phi \quad [4]$$

where $s_i = 2b\lambda_i$, for i=1,2,3.

First, we consider the trivial case of isotropic tensors, i.e., $\lambda_i = \lambda$. The exponent in Eq.[4] can be factored out as a scalar multiplier, and the result is the sum of an infinite number of sampling orientations which Batchelor et al.[4] have previously shown takes the form in Eq. [2]. For a more general case, however, the evaluation of Eq.[4] becomes more complex. Space prohibits detailed working, (utilizing Mathematica), but the end result is:

$$\langle f(l,m,n,s_1,s_2,s_3)\rangle = \frac{b^2}{2\sqrt{\pi}} e^{-s_3} \frac{\Gamma(1+l+m)\Gamma(\frac{1}{2}+n)}{\Gamma(\frac{3}{2}+l+m+n)} \sum_{k=0}^{\infty} \left(\frac{(1+l+m)_k}{k!(\frac{3}{2}+l+m+n)_k} \sum_{q=0}^{k} \left(\binom{k}{q} (s_3-s_1)^{k-q}(s_1-s_2)^q \frac{\Gamma(\frac{1}{2}+m+q)}{\Gamma(1+m+q)} {}_2F_1(-2l;\tfrac{1}{2}+m+q;1+2m+2q;2) \right) \right) \quad [5]$$

where ${}_1\tilde{F}_1(a;b;z) = {}_1F_1(a;b;z)/\Gamma(b)$ is the regularized confluent hypergeometric function, ${}_1F_1(a;b;z) = \sum_{k=0}^{\infty} \left((a)_k z^k / (b)_k k!\right)$ is the Kummer hypergeometric function, and $(a)_n = a(a+1)(a+2)\cdots(a+(n-1))$ is the Pochhammer function. We now assume a general tensor such that (s_1, s_2, s_3)=($2b\lambda_1$, $2b\lambda_2$, $2b\lambda_3$)=(0.1, 0.01, 0.001) and evaluate the leading three diagonal elements of M', ($M_{1,1}$, $M_{2,2}$ and $M_{3,3}$) using Eq. [5]. The results are f(2,0,0, 0.1, 0.01, 0.001) = 0.195056…; f(0,2,0,0.1,0.01,0.001) = 0.195754… and f(0,0,2,0.1,0.01,0.001) = 0.190843…. This indicates that the leading terms are not equal and therefore the prescription given in Eq. [2] is violated.

CONCLUSION: In the limit of an infinite number of sampling vectors, the precision matrix is rotationally invariant *only* for the trivial case of isotropic tensors – a result previously outlined elsewhere[5]. Even in this infinite limit, however, for a given anisotropic tensor, the leading 3 diagonal terms of M' are unequal and hence the condition laid out in Eq. [2] is violated. As discussed earlier, just one example of non-conformance to Eq. [2] is sufficient to conclude that, despite claims by some groups to the contrary, it is indeed NOT possible to design a statistically rotationally invariant sampling scheme.

REFERENCES: 1. Basser et al. *TMI* 2003 ; 22 :785- ; **2.** Jones *MRM* 2004; 51:807-; **3.** Skare et al. *J Magn Reson* 2000; 147: 340- ; **4.** Batchelor et al. *MRM* 2003; 49:1143-; **5.** Jones *Proc ISMRM* 2003, p. 2118. **6.** Koay et al. *J Magn Reson* 2006; 182:115-; **7.** Hext *Biometrika* 1963; 50:353-;